Astronomy & Astrophysics manuscript no. Paper
September 6, 2012

© ESO 2012

arXiv:1209.0216v2 [astro-ph.GA] 4 Sep 2012

A multi-waveband study of the Magellanic Bridge

G. Bagheri[1], M.-R.L. Cioni[1,2,*] and R. Napiwotzki[1]

[1] Centre for Astrophysics Research, College Lane, Hatfield AL10 9AB, United Kingdom
[2] University Observatory Munich, Scheinerstrasse 1, 81679 München, Germany

Received 2012 October 11/ Accepted XXX

ABSTRACT

Context. The Magellanic system comprises the Large Magellanic Cloud (LMC), the Small Magellanic Cloud (SMC), and the less frequently observed Magellanic Bridge and Magellanic Stream. The Bridge is traced by neutral gas and has an observed stellar component, while the Stream consists of gas only, with no observed stellar counterpart to date.
Aims. This study uses previously unexplored infrared observations from the 2MASS and DENIS surveys and optical observations from the SuperCOSMOS survey to investigate the stellar content of the Magellanic Bridge.
Methods. Catalogues were created of the 2MASS, DENIS and SuperCOSMOS surveys in the direction of the Bridge and density maps, colour-magnitude and two colour diagrams were analysed. A study was also carried out on removing the Galactic foreground population in the direction of the Magellanic Bridge which was an important consideration due to the low stellar density within the Bridge.
Results. This study finds that the stellar density varies in the Bridge region with a north–south division, and that the Magellanic Bridge comprises of an old stellar population in addition to the younger population already known.
Conclusions. The formation of the Magellanic Bridge is likely to have occurred from a tidal event between the LMC and SMC drawing most of the material into it from the SMC. An older population in the Bridge indicates that a stellar content was drawn in during its formation together with a gas component. Specific attention was given to the removal of the Galactic foreground due to the low stellar density within the Bridge.

Key words. Galaxies: interactions, Galaxies: individual, Galaxies: stellar content

1. Introduction

The Magellanic Clouds are irregular dwarf galaxies within our Local Group at a distance of $50-60$ kpc from the Milky Way (Cioni et al. 2000). The Clouds consist of the Large Magellanic Cloud (LMC) and the Small Magellanic Cloud (SMC) which are connected by the Magellanic Bridge. The system also contains the Magellanic Stream which is a gaseous feature between the Clouds. It is widely believed that the Bridge was formed tidally (Mathewson 1985) while the formation of the Stream is still under debate as a tidal feature or a trailing creation of the Clouds from ram pressure from a low density ionized halo (Mastropietro et al. 2004). The Magellanic Clouds are a prime template for studying galaxy interactions, the derivation of their star formation history (SFH), and the interaction between the Clouds and the Galaxy (Cioni et al. 2011).

The Magellanic Bridge spanning $\sim$ 13 kpc contains a known stellar population (Irwin et al. 1990). Star formation after a tidal event can be studied by examining the ages, chemical abundances and kinematics of the Bridge population. To date, only young populations have been observed in the Bridge region, this implies that at the time of the tidal interaction, only gas was drawn out of the Clouds to create the Bridge, and that the inter-cloud population of the Bridge formed some time afterwards. If an older population were to be observed, it could imply that stars as well as gas were stripped from the LMC and SMC in the event that formed the Bridge. As tidal interactions should affect both stars and gas, an older population is expected within the Bridge.

This study addresses the inter-cloud population in the Magellanic Bridge and its distribution. An overview of the Magellanic Bridge is presented in Sect. 2. Section 3 describes the 2MASS, DENIS and SuperCOSMOS data used in this study and their analysis. A study of the stellar densities in the Bridge is carried out in Sect. 4.1. The results of the data analysis, stellar density maps and colour-magnitude diagrams (CMDs), are shown in Sect. 4.2. A discussion is given in Sect. 5 and the conclusions are summarised in Sect. 6.

2. The Magellanic Bridge

Hindman et al. (1963b), carried out observations and digital reductions for a survey of neutral hydrogen (HI) in the direction of the Magellanic Clouds and noted a considerable amount of gas between the two Clouds. In a follow-up paper Hindman et al. (1963a) concluded that the Magellanic Clouds were encompassed by a halo of gas. They also observed double peaks in the velocity of the SMC gas possibly indicating two separate masses one behind the other. Deinzer & Schmidt (1973) used observations of the 21cm H I line to show a physical connection of gas between the Clouds. This has been subsequently confirmed by other authors and Irwin et al. (1990) showed that the SMC Wing extends towards the LMC into the Bridge. This is supported by the observation of several regions within the

---

Send offprint requests to: g.bagheri@herts.ac.uk

[*] Research Fellow of the Alexander von Humboldt Foundation

Bridge containing young blue stars. Another investigation by Grondin et al. (1991), using CCD photometry of three regions within the Bridge, indicated a population of young stars approximately 10^8 years old. Grondin et al. (1992) observed two regions in the SMC wing and one in the centre of the Bridge. They found that the SMC Wing population has a large line of sight depth of 12 ± 8 kpc and that some associations are as young as 16 Myr. This study also observed that the inter-cloud stars were younger and less numerous than the LMC, their distribution was not continuous, and were likely to have formed from SMC gas.

Mathewson (1985) proposed that the SMC had a close encounter with the LMC warping the LMC disk and forming the Bridge while tidally fissioning the SMC in a process of irreversible disintegration. It was also postulated that the Clouds were not bound to the Galaxy and were approaching from the direction of Andromeda. Grondin et al. (1991) proposed a scenario where a close encounter of the LMC and SMC occurred, triggering star formation in the densest parts of the inter-cloud region leaving dense gas on the SMC side. Their analysis confirmed that the Bridge associations on the west side formed first. It was then suggested by Grondin et al. (1992) that a tidal interaction between the Clouds was on a milder scale than seen elsewhere in external galaxies but caused the formation of a disk around the SMC which was then stripped to form the Bridge. A ring-like feature around the SMC has been suggested by Cioni et al. (2006) and also Harris & Zaritsky (2004) corresponding to an older population of about 2.5 Gyr.

Authors such as Putman & Gibson (1999) and Muller et al. (2004) have supported that the Bridge may have been formed from a close encounter 200 Myr ago between the LMC and SMC, but this is challenged with ideas that the Bridge was formed later due to the difference between the Bridge and SMC populations and that the ages of associations in the Bridge are closer to 6 Myr. It is possible that the Bridge was formed in a close encounter 200 Myr ago but that star formation was triggered by another event about 6 Myr ago.

Muller et al. (2003) showed that the northern and southern parts of the Magellanic Bridge represent two distinct arms of gas, emanating from the SMC, which challenges previous theories that the Bridge was formed of a single filamentary feature. The southern part of the Bridge has more turbulent H I similar to the SMC while the northern part is more tenuous. This study is followed by Muller & Bekki (2007) who also found that there exists velocity bi-modality within the Bridge and that large HI loops are present up to 1.3 kpc in diameter spanning across the northern and southern parts of the Bridge. These authors suggest that the northern part of the Bridge gas is a radially extended arm of the SMC and is not connected to the LMC but that the southern part was drawn from the SMC in a LMC–SMC interaction. There are many authors that argue that the Bridge is predominantly composed of SMC matter. Hambly et al. (1994) using the 3.9m Anglo-Australian telescope Echelle spectrograph, observed one main sequence B-type star and one B-type supergiant and determined the chemical composition and spin temperature of the interstellar medium to determine the temperature structure within the Bridge. They find that the stars had a mild He deficiency of $0.1 - 0.2$ dex and a heavy element deficit of $0.5 - 1$ dex with respect to the Galaxy which both reflect the compositions within the SMC and not the LMC.

This may aid in the confirmation of Bridge formation scenarios that the Bridge formed from a tidal interaction between the Clouds 200 Myr ago stripping gas from the SMC and that an event around 6 Myr ago triggered star formation from molecular condensation with the presence of CO (Muller et al. 2003). It is also possible that the northern part of the Bridge, not connected to the LMC, may have been stripped from a different region of the SMC that contained stars at the time of the stripping event. In order to confirm this, an older population will have to be observed in the Bridge which to date has not been found. Harris (2007) used the MOSAIC II camera on the 4m telescope at CTIO to observe twelve regions in the Magellanic Bridge searching for an old stellar population which may have been stripped from the SMC. The spatial distribution of these regions follows the H I gas ridge line in the Bridge and extends to right ascension $\sim 3^{\rm h}$ where the H I surface density falls to below the critical threshold for star formation as determined by Kennicutt (1989). Harris work showed no detection of an old population in the Bridge indicating perhaps that the stars were all formed in-situ.

3. Data

The aim of this work is to establish source catalogues in the region of the Magellanic Bridge prior to the analysis of observations using the VISTA telescope (Emerson et al. 2004) as part of the VISTA survey of the Magellanic Cloud system (VMC; Cioni et al. (2011)). As such, the main source of data used in this work is near-infrared (NIR) photometry from the Two Micron All Sky Survey (2MASS) and the Deep Near-Infrared survey of the southern sky (DENIS) respectively. Optical SuperCOSMOS data are also used for comparison. The 2MASS, DENIS and SuperCOSMOS catalogues are publicly available. The Bridge area was chosen from previous definitions of the Bridge coverage by Kerr & de Vaucouleurs (1955), McGee & Newton (1981) and Harris (2007). This corresponds to an area between the LMC and SMC from 20° to 70° in right ascension (α) and from $-69°$ to $-77°$ in declination (δ).

DENIS is a complete survey in one optical band I (0.8μm) and two NIR bands, J (1.25μm) and $K_{\rm s}$ (2.16μm) of the southern sky (Epchtein 1994). The limiting magnitudes of the survey are 18.5, 16.5 and 14.0 mag in I, J and $K_{\rm s}$ bands, respectively with a 1σ photometric uncertainty of < 0.1 mag and a pixel size of $1''$ in I and $1''$ in J and $K_{\rm s}$ citepmr00a. From this catalogue there were $\sim 170,000$ sources extracted in the Bridge area. [1]

2MASS collected raw imaging data covering 99.998% of the celestial sphere in the NIR at J (1.25μm), H (1.65μm) and $K_{\rm s}$ (2.16μm) between 1997 and 2001 (Skrutskie et al. 2006). The point source detection level is 15.8, 15.1, and 14.3 mag at the J, H, and $K_{\rm s}$ bands, respectively, for the vast majority of observations. The 1σ photometric uncertainty is < 0.03 mag for bright sources with a pixel size of $2.0''$. From this catalogue there were $\sim 300,000$ sources extracted in the Bridge area. [2]

For completeness in the analysis of the Magellanic Bridge, a comparison of the NIR data above was done with the SuperCOSMOS catalogue which covers this region with reasonable depth. SuperCOSMOS is an advanced photographic plate digitising machine which digitised sky survey plates taken with the UK Schmidt telescope (UKST), the ESO Schmidt, and the Palomar Schmidt telescopes for the entire sky. The pixel size of the SuperCOSMOS surveys is 0.7″. The SuperCOSMOS Sky Surveys are beneficial due to the small pixel size and the availability of proper motions and variability information. For this work the UKST southern Survey catalogue was used with band pass filters B (0.45μm), R (0.66μm) and I (0.8μm) and limiting magnitudes of $B = 22.5$, $R = 20.0$ and $I = 19.5$ mag (Hambly et al. 2001) with photometric errors of 0.3-0.5 mag. From this catalogue $\sim 450,000$ sources were extracted in the Bridge area. [3]

---

[1] http://irsa.ipac.caltech.edu/Missions/denis.html

[2] http://irsa.ipac.caltech.edu/Missions/2mass.html

The data for the Bridge region was downloaded from the 2MASS, DENIS and SuperCOSMOS web interfaces using columns of α, δ, magnitudes and photometric uncertainties with different flags used to only extract stellar objects. In 2MASS flags were used to remove objects with JHK artefact contamination and/or confusion and contamination from extended sources. DENIS flags were applied to remove multiple observations of sources and maintain image and source quality. Catalogues were then created by selecting only stellar objects observed within each survey. The results in this work are mostly affected by the photometric uncertainties in the magnitudes. To reduce the photometric uncertainties near the sensitivity limits, cuts at $K_{\rm s} = 14$ and $R = 17$ mag were applied. The photometric uncertainties of colours are derived from $\sigma_{\rm A-B} = \sqrt{\sigma^2_{\rm A} + \sigma^2_{\rm B}}$. In order to complete a full study of the Magellanic Bridge the catalogues of 2MASS and DENIS were combined to form a catalogue with I, J, H and $K_{\rm s}$ bands. Because the 2MASS catalogue reaches fainter magnitudes than DENIS in the J and $K_{\rm s}$ band, and the photometric errors are smaller in 2MASS (Fig. A.1), after cross matching the catalogues, the J and $K_{\rm s}$ bands of DENIS were discarded. The 2MASS and DENIS samples were matched using coordinates and a separation of 0.5″ to ensure that the stars in the combined catalogue were the same stars in each catalogue. From the original $\sim 170,000$ objects in DENIS and $\sim 300,000$ in 2MASS, the combined catalogue contained $149,290$ objects observed in both 2MASS and DENIS.

Table 1. Number of stellar objects in each catalogue of the Magellanic Bridge

Catalogue	Filter	Number of sources
2MASS	J, H,$K_{\rm s}$	$\sim 300,000$
DENIS	I, J, K	$\sim 150,000$
2MASS-DENIS	I, J, H, $K_{\rm s}$	$\sim 150,000$
SuperCOSMOS	B, R, I	$\sim 450,000$

Table 1 contains the different catalogues used in this work with their wave bands and also the number of sources in each catalogue. The inclusion of all three surveys is a useful tool in analysing the Bridge population especially in terms of a comparison between red and blue objects.

4. Results

The results of this work look at the structure of the Bridge, and also the populations within. The work on the structure, makes use of a statistical removal of the Galactic foreground from the 2MASS and SuperCOSMOS catalogues. The study into the stellar populations uses a more robust method of foreground removal and also makes use of the DENIS catalogue combined with 2MASS.

4.1. Density within the Bridge

In order to investigate the density of the stellar population within the central Bridge (30-60 deg in RA, to avoid too greater LMC and SMC contamination), stellar density maps were created from the catalogues of 2MASS and SuperCOSMOS prior to the consideration of a Galactic foreground population (Fig. 1), to look into different stellar densities contrary to the particular populations. Since a mostly structureless, smooth spacial distribution of foreground stars and background galaxies can be expected across the Magellanic Clouds, any structure found should reflect the Bridge population.

The positions of the stars in the 2MASS and SuperCOSMOS catalogues were binned, and smoothed number density maps were created with bin size dependent on the total number of stars present. Bins were optimized at $\Delta\alpha = \Delta\delta = 0.2^\circ$ as this was the bin size that produced the clearest contours on the density maps with an average of 8 and 41 objects in each bin for the 2MASS and SuperCOSMOS catalogues respectively.

Regions previously observed by other authors were placed (not to scale) on top of the maps. The most significant observations in this region were those by Harris (2007) using tiles of $36' \times 36'$ who looked at different regions across the Bridge. The position of the VMC tiles (1.5×1 deg) is also indicated on the maps to give an idea of the larger region that VMC will cover in comparison to previous observations. The exact location of the VMC tiles has been decided according to the present study, with tiles placed at higher declinations than initially planned with the expectation to observe different stellar populations. The 10 sigma sensitivity (Vega) in Y, J and $K_{\rm s}$ are 21.1, 21.3 and 20.7 respectively. Please refer to Cioni et al. (2011) for details on the VMC observations.

It can be seen from the 2MASS density map of the Bridge (Fig. 1), that the stellar distribution is not even. The Harris tiles (medium size boxes) follow the H I ridge line within the Bridge and the stellar populations are distributed along it, but also in other regions. The SuperCOSMOS map (Fig. 1) shows an apparent increase in star numbers in the northern part of the Bridge just outside the regions observed by Harris. One would expect to find a greater number of young blue objects in the SuperCOSMOS catalogue compared to the redder objects in the 2MASS catalogues. Therefore the greater number of objects in the northern part of the Bridge in SuperCOSMOS opposed to 2MASS indicates the possibility of a denser younger population in the north of the Bridge. It also appears from the 2MASS density map that the number of objects increases towards the south, this could indicate a possible difference in young and older populations from the north to the south.

---

[3] http://www-wfau.roe.ac.uk/sss/



Fig. 1. Stellar density maps of the Magellanic Bridge from the 2MASS (left) and SuperCOSMOS (right) catalogues, with bin size of 0.2×0.2 deg^2. The rectangles represent the locations (not to scale) of VMC observations (red) (Cioni et al. 2011), and fields observed by Harris (2007) (blue). Contour levels are at: 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60 and 80 stars. North is at the top and east is to the right with the LMC to the left and SMC to the right
.

In order to confirm that the observed stellar density fluctuations in the Bridge are truly a feature of the Bridge and not from the Galactic foreground population a study into the Galactic foreground was carried out. A target region was chosen to represent a purely Galactic foreground population in the direction opposite to the Magellanic Clouds across the Galactic plane, at the same Galactic longitude but at positive Galactic latitude. This region was chosen as it represents a similar location to the Clouds in terms of reddening effects and foreground population, but without the galaxies dominating the region it better represents a purely Galactic population. For this work, catalogues were downloaded at (292°, 39°), to cover an area equivalent to that of the central Bridge region ($\sim 7° \times 8°$). Flags were used to clean up the catalogues as in Sect. 3 and density maps were created following the same procedure as was applied to the Bridge field to identify the nature of the stellar density.

It can be seen from the stellar density maps of the foreground field for 2MASS and SuperCOSMOS (Fig. 2) that in each case, with the same binning as the Bridge regions, the stellar density across the foreground region is not entirely smooth. A preliminary observation is that the average number of stars in the Bridge catalogues is larger than that in the foreground. In the SuperCOSMOS map there is a visible gap due to the photographic plates used to obtain the photometry. The overall stellar densities allow the preliminary conclusion, that a significant fraction of the 2MASS and SuperCOSMOS objects are stars from the Bridge population. However, interpretation of these maps is dampened by the contribution of the general Galactic foreground ($\sim 50\%$ and $\sim 80\%$ in the SuperCOSMOS and 2MASS maps, respectively). Both Bridge maps display some spacial structure. However, some structure can also be seen in the foreground. In order to clarify whether true Bridge structure remains, a statistical removal of the Galactic foreground was applied as follows.

In order to statistically remove Galactic foreground objects from the Bridge population, the foreground field was binned in colour-magnitude space to obtain a number of objects in each bin of colour and magnitude. For the 2MASS catalogue, the field was binned in $J - K_{\mathrm{s}}$ vs K_{s} and in SuperCOSMOS the binning was in $B - I$ vs I. In each case the colour and magnitude bins were 0.2 and 0.5 in size respectively for 2MASS, and 0.5 and 0.5 for SuperCOSMOS. This process was repeated for the Bridge region with the same binning for each catalogue.

The next step in the removal process was to calculate the probability for each bin, of an object belonging to either the Bridge or the foreground. This was calculated by comparing the number of Bridge and foreground objects in each bin, with respect to the total number of objects in each bin. Once a probability had been assigned to each bin in colour-magnitude space, a cut was made to select objects with a probability of greater than 0.20 of belonging to the Bridge, to avoids noise and objects with biases. The objects remaining were then analysed to seek density fluctuations that are genuinely Bridge features and not an effect of the Galactic foreground population.

Histograms were created for the Bridge and Galactic foreground fields in the 2MASS and SuperCOSMOS catalogues (Figs. 3, 4, top). A Poisson fit was applied to each of the data sets representing the average number of objects in each catalogue to identify any fluctuations that are not expected statistically. As well as for the full catalogues, histograms were created after the statistical removal of Galactic foreground, for objects that are likely to be Bridge stars in the NIR and optical catalogues, as



Fig. 2. Stellar density maps of the Galactic foreground field from the 2MASS (left) and SuperCOSMOS (right) catalogues, with bin size of 0.2×0.2 deg^2. Contour levels are at: 10, 15 and 20 stars. North is at the top and east is to the right with the LMC to the left and SMC to the right.



Fig. 3. Histograms of no. of stars in a given bin range for the Bridge in the 2MASS catalogue (red) and foreground field (blue, top) in 2MASS, and also the 2MASS catalogue of the Bridge after the statistical removal of the foreground (bottom). A Poisson distribution was computed for each figure representing the expected distribution on the average number of stars per bin of each catalogue
.



Fig. 4. Histograms of no. of stars in a given bin range for the Bridge in the SuperCOSMOS catalogue (red) and foreground field (blue, top), and also the SuperCOSMOS catalogue of the Bridge after the statistical removal of the foreground (bottom). A Poisson distribution was computed for each figure representing the expected distribution on the average number of stars per bin of each catalogue
.

well as objects likely to make up the Galactic foreground in the direction of the Bridge (Figs. 3, 4, bottom). It should be noted that the Poisson fit for the SuperCOSMOS foreground field is slightly to the left due to the small gap in SuperCOSMOS observations as seen in Fig. 2. This effect is compensated for in the removal of the foreground from the Bridge by extrapolating the rest of the field to fill this gap.

It can be seen from Figs. 3 and 4, that prior to and after the removal of the foreground objects, there remains a greater number of Bridge objects in the catalogue which are likely to be real Bridge stars and are above the expected number given by the Poisson distribution. The statistical foreground removal is based on the positions of the objects in colour-magnitude space, indicating that the larger number of objects prior to removal are of different populations to those dominating the Bridge. A further study of the NIR catalogue in the direction of the Bridge is carried out in Sect. 4.2, using a more robust method of removing the Galactic foreground objects, to identify new, older populations within the Bridge.



Fig. 5. Stellar density maps of the Magellanic Bridge from the 2MASS (left) and SuperCOSMOS (right) catalogues after the statistical removal of the Galactic foreground multiplied by the probability of a star belonging to the Bridge, with bin size of 0.2×0.2 deg^2. Contour levels are at: 60, 70, 80, 90, 100, 120, 140, 160, 180 and 200 stars. North is at the top and east is to the right with the LMC to the left and SMC to the right

.



Fig. 6. Number of stars in bins of Dec for the central Magellanic Bridge ($\alpha = 30° - 60°$) in the SuperCOSMOS (crosses) and 2MASS catalogue (squares, scaled by a factor of 3) after the statistical removal of Galactic foreground. The Bridge is binned over all RA with a bin size of 0.5° in Dec. Vertical lines represent the range of Dec of Harris' (2007) observations. Error bars are included but are smaller than the symbols

.

Fig. 5 shows stellar density maps of the Bridge in 2MASS and SuperCOSMOS after the statistical removal of the Galactic foreground. It can be seen from these maps that the two regions of over-density in the SuperCOSMOS catalogue (Fig. 1) still remain after the removal of foreground. The SuperCOSMOS catalogue also shows a distinction between the northern and southern parts of the Bridge with a denser population in the north. The 2MASS map shows a lower density of objects overall, but also shows regions of over-density remaining after the foreground removal.

Fig. 6 shows that the highest density of blue objects is not along the main concentration of H I gas around which Harris (2007) carried out his observations of Bridge objects. This work shows a peak in stellar density of blue objects in the northern part of the Bridge decreasing towards the south. Harris does however state that due to ram pressure from the Magellanic Clouds passing the Milky Way, then tidally stripped stars can be displaced from the main gas concentration, indicating that perhaps there has been movement in the position of the main concentration of H I gas.

4.2. Stellar populations

The purpose of this section is to identify an older population in the Bridge using NIR observations. In order to analyse stellar populations, foreground Galactic stars towards low density regions of the Magellanic System, like the Bridge, represent a non-negligible source of contamination as Galactic objects will appear with similar colours and magnitudes as Bridge objects, and will make up most of the objects observed. The foreground population was removed here using CMDs and two colour diagrams. Magnitude cuts near the sensitivity limits of the surveys were also applied. Two offset regions were chosen, one above and one below the Bridge to obtain a sample of the foreground population. The offset fields were chosen to be 2° × 2° in size centred at α = 3:26:00, δ = −59:53:0 and α = 4:01:31.91, δ = −80:58:19.8. The offset fields were analysed for Galactic dust reddening by examining dust extinction maps (Schlegel et al. 1998) as these fields are assumed to comprise of only Galactic foreground stars. The average absorptions ($A_I = 0.073$, $A_J = 0.033$, $A_H = 0.021$, $A_K = 0.014$) are sufficiently low to allow use of these fields in the removal of Galactic foreground stars from the Magellanic Bridge population and this is re-enforced as the Magellanic Clouds lie sufficiently above the Galactic plane for reddening effects to be low and therefore no correction for reddening has been made. Two Bridge fields of 2° × 2° were also chosen to represent the Bridge population, in a test for the foreground removal process, in the direction of the highest stellar density at α = 2:00:00, δ = −75:30:00 and α = 2:20:00, δ = −70:00:00. It should be noted that the reddening effects of the Bridge were also examined and the average absorptions ($A_I = 0.094$, $A_J = 0.043$, $A_H = 0.028$, $A_K = 0.018$) are also low enough here and similar to the offset fields, to continue without a reddening correction for the Bridge region.

The two offset and Bridge fields were then compared using a range of CMDs and two colour diagrams to find the best distinctions between the Bridge and the offset field stars. In this work, the DENIS catalogue was also included to make use of the I band for the more robust method of removing Galactic foreground. The foreground was removed from a combined 2MASS-DENIS catalogue. A CMD was also used to make cuts in the K_s magnitude for the combined catalogue near the sensitivity and brightness limits to reduce statistical fluctuations in the data (Fig. A.1). These cuts were taken at $K_\mathrm{s} = 9.0$ and $K_\mathrm{s} = 13.8$ mag for the bright and sensitivity limit respectively. This analysis was repeated for all bands and was used in this work to determine the point at which the photometric uncertainty was too large for the data to be reliable. Figure A.1 shows that the photometric uncertainty increases for stars brighter than $K_\mathrm{s} \sim 8.5$ mag due to special processing of bright sources in 2MASS, and a cut around this level excludes Galactic giant stars as they will appear as very bright objects. The cut at $K_\mathrm{s} = 13.8$ was chosen as a compromise between the higher photometric uncertainties near the sensitivity limit of the catalogue and not removing faint objects in the Magellanic Bridge. Our method for the removal of foreground stars makes primarily use of the different IR colours of late type dwarfs and giants (Bessell & Brett 1988).



Fig. 7. $I - K_{\mathrm{s}}$ histogram for the Bridge field of the 2MASS-DENIS combined catalogue showing a drop in stellar density at the chosen cut of 2.3
.

The best separation of the Bridge and Galactic populations was observed in the $I - K_{\mathrm{s}}$ vs K_{s} hess diagram (Fig. 7). A value of $I - K_{\mathrm{s}} > 2.3$ was initially taken by eye as the best cut to separate the Galactic foreground from the Bridge population. Note that $J - K_{\mathrm{s}}$ vs K_{s} was not included in the cuts as this combination of colour and magnitude are used below to identify the stellar populations and as such were not used to filter out objects.

Photometric errors of the photometry cause a scattering of foreground stars into clean regions in colour space. Our aim was to find an optimal trade off between removing as much Galactic foreground contamination as possible without removing too many genuine Bridge stars. For this purpose we determined the best S/N value using the formula $\mathrm{S/N} = \frac{N_{\mathrm{Bridge}}}{\sqrt{N_{\mathrm{Bridge}} + N_{\mathrm{Foreground}}}}$ for the total number of Bridge stars and total number of Galactic foreground stars remaining after a cut in each colour combination. This application, with an optimum S/N of 10 produced a final Galactic foreground cut of $I - K_{\mathrm{s}} > 2.3$ mag.

Figure 8 shows the two colour diagram $H - K_{\mathrm{s}}$ vs $J - H$ of the combined 2MASS-DENIS catalogue and the Galactic dwarf and giant sequences (Bessell & Brett 1988) which aided the Galactic foreground removal process. From this comparison a cut at $J - H = 0.70$ was chosen initially by eye to remove the Galactic dwarfs from the catalogue which will cause contamination to the Magellanic objects. Again a range of values around this $J - H$ cut were checked for the best S/N ratio (7.9) resulting in a value of $J - H > 0.75$ mag. The remaining objects were then further selected in $H - K_{\mathrm{s}}$ (S/N = 13) to remove any remaining Galactic population by making a cut at the point in which the two population arms begin to separate, i.e. the Magellanic giant population begins to divert away from the main bulk of Galactic M dwarf objects. The final colour selection criteria for the 2MASS-DENIS combined catalogue were $I - K_{\mathrm{s}} > 2.1$, $J - H > 0.75$ and $H - K_{\mathrm{s}} > 0.32$ mag which were applied together with the magnitude cuts at $K_{\mathrm{s}} = 9.0$ and $K_{\mathrm{s}} = 13.8$ mag.

A similar method of the removal of Galactic foreground objects was carried out for the optical SuperCOSMOS data using the same Bridge and Galactic foreground fields as with the combined 2MASS-DENIS catalogue. These cuts were tailored for the identification of a young blue population. Magnitude cuts in the I band were applied to account for the brightness and sensitivity limits of the catalogue, taken at $I = 18.4$ and $I = 10$ at a similar level with respect to those in the combined 2MASS-DENIS catalogue. A CMD, $B - I$ vs I, and two colour diagram, $R - I$ vs $B - R$, were used to separate the Galactic foreground sources from the Magellanic sources. The cuts in these diagrams were initially taken at $B - I < 0.4$, $R - I < 0.35$ and $B - R < 0.5$ mag. After checking the spread around these values the values of $B - I < 0.5$, $R - I < 0.3$ and $B - R < 0.6$ mag were chosen (with optimum S/N = 24.4, 24.7 and 25.1 respectively) for the final Galactic foreground removal. A cut at the blue end of the catalogue was also applied to remove background galaxies at $B - R > -0.5$ mag. This cut was confirmed to be accurate as all sources with $B - R < -0.5$ mag are galaxies according to the SIMBAD[4] astronomical database. Note that these selection criteria remove Magellanic Cloud red giants as well.

The SuperCOSMOS objects have been compared to the colours of known main sequence and G-M giant objects in Fig. 10. A few runaway Galactic stars and hot evolved objects (horizontal branch, post-Asymptotic giant branch (AGB) and white dwarf stars) may slip through but numbers are small as confirmed by the CMD of the foreground fields. It can be seen here that although originally a statistical process, the method of Galactic foreground removal in the SuperCOSMOS catalogue excludes Galactic main sequence stars and giants.

The number of stars in the Magellanic Bridge area is reduced to 449 in the combined 2MASS-DENIS catalogue and 274 in the SuperCOSMOS catalogue after the Galactic foreground removal.

A further extension to the Galactic foreground removal process used SIMBAD over the entire Bridge region to investigate the candidate Bridge objects in the 2MASS-DENIS combined catalogue and the SuperCOSMOS catalogue with a breakdown of objects in Table 2. Using this method a further 74 objects were removed from the 2MASS-DENIS combined catalogue as background galaxies and 7 objects were removed from the SuperCOSMOS catalogue as Galactic stars and background galaxies representing ∼ 2.6% and ∼ 17% of the final catalogues. It should be noted that no previous attempt was made to remove background galaxies from the 2MASS-DENIS catalogue beyond using the point source flags. The remaining 365 objects in the 2MASS-DENIS combined catalogue and 267 in the SuperCOSMOS catalogue are all likely to be objects with a Magellanic origin.

4.3. Colour Magnitude Diagrams

In order to characterise the stellar populations within the Bridge, a CMD wes created from the 2MASS-DENIS combined catalogue in $(J - K_{\mathrm{s}})$ vs K_{s}. Boxes are from Nikolaev & Weinberg (2000). These boxes were created to characterise the population of the LMC (Fig. 11), and have been adapted here for the Magellanic Bridge using the distance modulus at an average Bridge distance of 55 kpc. The boxes were then overlaid onto the observed CMD of the Bridge (Fig. 12).

There are twelve regions identified in Fig. 11 marked from A through L and several techniques were applied by Nikolaev & Weinberg to identify the populations in each region. The LMC populations were identified based on the

---

[4] http://simbad.u-strasbg.fr/simbad/



Fig. 8. $H-K$ vs $J-H$ two colour diagram for the 2MASS-DENIS combined catalogue (dots) and Bessell & Brett (1988) Galactic dwarfs and giants ((red) diamonds). Bold lines illustrate the foreground removal cuts at $H-K_s > 0.32$ and $J-H > 0.75$ mag
.

NIR photometry of known populations from the literature. The authors also performed a preliminary isochrone analysis, where they matched the features of the CMD with Girardi et al. (2000) isochrones, to derive the ages of populations and draw evolutionary connections among the CMD regions. The regions of the CMD present in the Bridge population from the 2MASS-DENIS combined catalogue after the removal of Galactic foreground (Fig. 13) are primarily E, F, J, D, K and L with a small contribution from I. Even considering photometric uncertainties in magnitudes and colour, the stellar population still fits well into the boxes characterising the different types of stars.

A summary of the types of stars found in the regions according to Nikolaev & Weinberg:

Region E covers the upper RGB and includes the tip of the RGB. The tip of the RGB is defined by the ignition of the degenerate helium burning core in old (low mass) stars, the helium flash (Renzini & Fusi Pecci 1988). Stars, after ignition, evolve rapidly to the horizontal branch.

Region F contains primarily oxygen-rich AGB stars of intermediate age that are the descendants of stars in region E. Some fraction of region F stars are long-period variables and reddened supergiants.

Region D is the most inhomogeneous part of the CMD due to the large colour range spanned. This region contains RGB and early AGB (E-AGB) stars. Region D also contains a significant fraction of AGB stars in transition from the E-AGB to the thermal pulsing AGB phase (TP-AGB), the stage at which the outer hydrogen shell is reignited (Iben & Renzini 1982). Most of the E-AGB stars in this region are M-type stars.

Region J sources are primarily carbon rich TP-AGB stars. These stars are descendants of oxygen-rich TP-AGBs in regions F and G. Region K contains extremely red objects identified as obscured AGB carbon-rich stars, as well as some dusty oxygen-rich AGB stars. Their large $(J-K_s)$ colours are due to dust in their circumstellar envelopes.

Stellar sources in region L could be reddened M giants, there would usually be a large contribution here from background Galaxies however after each source was located in the SIMBAD astronomical database known Galaxies were omitted from these results.

The population in region I is young, with an age of approximately 500 Myr. A significant contributor to the source density in this region are K- and M-type supergiants.

From the above population analysis, it can be seen that the youngest populations remaining within the Bridge after the removal of Galactic foreground are K and M giants with an age of $\sim$ 500 Myr. This age is older than the assumed age of the Bridge implying that the red objects here are all older than the Bridge.

4.4. An older and younger population

This section confirms and expands on the previously observed younger population within the Magellanic Bridge region, and also looks into the existence of a population older than the Bridge (200Myr) according to the standard scenario (Grondin et al. 1991), indicating that these stars



Fig. 9. $B-I$ vs $R-I$ Hess diagrams of the Bridge fields (top) and Galactic foreground fields (bottom) of the SuperCOSMOS catalogue showing the difference in populations. Bin size is 0.1×0.1 in colour. Contour levels are at: 2, 4, 6, 8 and 10 stars
.

were drawn into the Magellanic Bridge as stars, and did not form in-situ as was previously thought.

The remaining objects in the 2MASS-DENIS combined catalogue and the SuperCOSMOS catalogue after the removal of Galactic foreground were cross-matched to the SIMBAD database and were characterised in α and δ (Fig. 14). It can be seen here that the known LMC and SMC populations from the database are confined to each side of the Bridge in both the 2MASS-DENIS and SuperCOSMOS catalogues, with known blue Bridge objects in SuperCOSMOS spreading across the entire region. Objects categorised as other, such as IR and X-ray sources are evenly distributed within the Bridge. The objects of interest here are the unknown objects which in both catalogues make up for the majority of the populations and are spread across the Bridge.

The remaining objects in the 2MASS-DENIS and SuperCOSMOS catalogues were entered into the SIMBAD database and of previously published objects within this study, just 2.6% and 17% are classified as non-Magellanic Cloud objects. Applying this accuracy to the previously identified objects (objects that have also not been published), it can be expected that $\sim$ 228 of the unknown 2MASS-DENIS and $\sim$ 126 of the SuperCOSMOS objects belong to the Magellanic Clouds. Follow up observations, required to confirm the type and membership of these objects, will be presented in subsequent paper. Many of these previously unobserved objects reside in the northern part of the Bridge, which has been less explored than the region around the H I ridge in the south especially in the NIR.

Figure. 15 shows two CMDs comparing the population within a field of the Bridge (α = 2:00:00, $\delta = -75$:30:00) to an offset field (α = 3:26:00, $\delta = -59$:53:0) of the same size of $2^\circ \times 2^\circ$. The panel with the Bridge field also includes remaining blue objects in the northern part of the Bridge in the SuperCOSMOS catalogue after the removal of galactic foreground for comparison, with isochrones (Girardi et al. 2000) over-plotted representing objects with SMC metallicity (Z = 0.004) at an average Cloud distance of 55 kpc for a range of ages. It can be seen from this figure that the remaining blue northern objects are truly plausible Bridge B stars and not a manifestation of the foreground population or background galaxies. The blue objects with known classification are all on their main sequence phase or beyond. Many of the unclassified objects are less evolved stars on the main sequence. This result confirms the methods used here to find a remaining blue Bridge population outside of the regions studied by Harris (2007) and Demers & Battinelli (1998) who concentrated their studies along the H ridgeline at the central Bridge declination.

Table 2 shows a breakdown of the objects as identified by SIMBAD within the Magellanic Bridge for the red (2MASS-DENIS) and blue (SuperCOSMOS) objects respectively after the removal of Galactic foreground and background contamination. It should be noted that the objects identified as background Galaxies here were consequently removed from the final sample.

5. Discussion

The density maps of the Bridge region all display a structured distribution of stars. The SuperCOSMOS map shows that for the central region of the Bridge, the population has a differing pattern in the northern and southern parts. The 2MASS-DENIS density map displays a vertical ridge at the centre of the Bridge which could be an artefact of the DENIS data set (related to the vertical scanning of the sky) or a region of low density between two populations of stars perhaps one from the LMC and one from the SMC. This feature is not present in the 2MASS or SuperCOSMOS data suggesting it is simply due to a lack of data in the DENIS catalogue.

An interesting finding here is of the stellar distributions of blue and red objects within the Bridge. Firstly a new



Fig. 10. $B-R$ vs $R-I$ colour-colour plot of the SuperCOSMOS catalogue of the Magellanic Bridge with the colours of known MS stars and G-M giants ((red) diamonds). Bold lines represent foreground cuts at $B-R<0.6$ and $R-I<0.3$ mag
.



Fig. 11. Hess diagram, $(J-K_{\rm s})$ vs $K_{\rm s}$ of LMC stars from 2MASS observations. The boxes overlaid on the plot are taken from Nikolaev & Weinberg (2000) and the catalogues include foreground stars. Contour levels are at: 2, 4, 6, 8 and 10 stars
.



Fig. 12. Hess diagram, $(J - K_\mathrm{s})$ vs K_s, of the Magellanic Bridge from the 2MASS-DENIS catalogue. Boxes are as in Fig. 11 and foreground stars have not been removed from the catalogue. Here the boxes have been shifted by 0.25 mag to brighter magnitudes in K_s. Contour levels are at: 10, 20, 30, 40, 50 and 60 stars
.

Table 2. Different populations identified within the Magellanic Bridge from SIMBAD. [1] Irwin et al. (1990), [2] Cannon & Pickering (1918), [3] Hog et al. (1998), [4] Skrutskie et al. (2006), [5] Holmberg et al. (1974), [6] Abraham et al. (1995), [7] Westerlund & Glaspey (1971), [8] Demers & Irwin (1991).

Object	No. within 2MASS-DENIS	No. within SuperCOSMOS	Reference
Bridge star	0	63	[1]
SMC star	53	30	[1] [7]
LMC star	21	0	[1]
Magellanic Cloud object	4	11	[1] [8]
Background Galaxy	74	3	[4] [5]
Other known object	0	20	[2] [3] [6]
Unknown	275	129	

red population has been identified after the Galactic foreground removal. Also an analysis of the blue objects within the Bridge yields a very interesting result that indeed there is a widely spread young population as previously observed, but that this population extends further northward that previously observed and also that the density of the blue objects is higher in the northern Bridge than in the south and not mostly concentrated along the H I ridge as suggested by Harris (2007). This implies that ram pressure may have played a role in the Magellanic system as it has moved through the Milky Way halo displacing the stellar population within the Bridge.

The north-south divide observed here, agrees with the work of Muller et al. (2004) who showed that the Magellanic Bridge comprises two distinct arms of gas making up the northern and southern parts. His results on the gas are confirmed here from stellar density maps supporting the hypothesis that the Bridge is not formed of a single filamentary feature. A more turbulent southern arm and more tenuous northern arm does also explain the more tenuous, less populous northern part of the density maps with respect to the denser southern parts.

Harris (2007) created an infrared CMD of a 35 deg^2 region of the central part of the Magellanic Bridge compared to a region of the outer Bridge towards the LMC. He found that in the outer region of the Bridge there was an abundance of RGB and AGB stars associated with the LMC but in the central region there were hardly any of these populations. The work here uses a robust method of Galactic foreground removal and finds a population within the Bridge including RGB and AGB stars of $\sim$ 72% of the total Galactic foreground subtracted stellar Bridge population in the 2MASS-DENIS catalogue, greater than the number of these objects found by Harris. The ages of the RGB and AGB populations identified here could range from $1 - 5$ Gyr for redder, lower-luminosity stars



Fig. 13. CMD, $(J-K_\mathrm{s})$ vs K_s, of the central Magellanic Bridge region from the 2MASS-DENIS catalogue. Boxes are as in Fig. 11 and foreground stars have been removed from the catalogue. Here the boxes have been shifted by 0.25 mag to brighter magnitudes in K_s
.

(Frogel & Blanco 1990). This indicates the presence of a population older than the age of the Bridge that is likely to have been drawn in tidally at the time of the Bridge forming event. An alternative explanation could be that the Bridge is older than previously estimated. Olsen et al. (2011) have identified a population of SMC objects within the LMC. These objects have lower metallicity than the LMC but are consistent with SMC giants which supports the theory of tidal interaction.

Other recent studies (Chen et al. 2010) have used mid-IR Spitzer IRAC and MIPS observations to detect a number of young stellar objects (YSOs) in the Magellanic Bridge, mostly nearby high H I density regions. They have also studied how star formation has proceeded and estimated the star formation efficiency in the Bridge by looking for Herbig Ae/Be stars using known catalogues of blue stars and stellar clusters. If the new populations in this work are genuine Bridge members they will contribute to the present knowledge of star formation within this region.

The Magellanic Bridge is one tidal Bridge between two irregular galaxies. Bekki (2008) investigated a Bridge that formed tidally between the outer halos of M31 and M33 made only from H I gas with no stars associated to it. Its origin is also under debate and it could be in-fact a tidal stream like the Magellanic Stream containing no or a very small stellar component. Smith & Redenbaugh (2010) have used the GALEX ultraviolet telescope to study various tidal features including bridges, spiral arms and tails of gas. There are many tidal features in existence, some with and some without stars within the gas in galaxies out to distances of 50 kpc (Pakzad et al. 2004). A more detailed study is required to determine a model to distinguish a tidal bridge from a stream, spiral arm or tail and a study into the conditions that form these features will give more information to the likely presence of stars within them. The Magellanic Bridge has had a known stellar population however, in the Magellanic Stream no stars have been detected within the H I gas. Perhaps unless drawn in at formation, tidal features do not contain a stellar population until an event occurs that is strong enough to trigger star formation within the tidal feature in an environment with a high enough gas density. Further observations of tidal systems, including the Magellanic Clouds with VISTA, are needed to give us a better picture of the interactions that form these features and star formation within.

6. Conclusions

This is the first infrared study of the Magellanic Bridge that confirms previous radio and optical studies which found that:

1. the Bridge contains a stellar population that is not smooth and consists of clumpy regions,
2. both blue and red stellar populations are way more extended than the region of high H I density,
3. the highest numbers of blue stars are in the northern Bridge region, different from expectations based on the morphology of the H I gas,
4. there is a north-south divide visible in the stellar population of the Bridge.

The above indicates that there may be an older population within the Bridge that has not yet been observed as the majority of current observations have been restricted



Fig. 14. The spatial distribution of the different populations present within the Magellanic Bridge from the 2MASS-DENIS catalogue (top) and the SuperCOSMOS blue catalogue (bottom) after the removal of Galactic foreground and background galaxies. The different symbols represent: known 2MASS-DENIS LMC sources or SuperCOSMOS known Bridge objects (squares), known SMC sources (stars), known Magellanic Cloud halo sources (diamonds), other known sources (triangles) and new sources (plusses) from SIMBAD
.

to the H I ridge line and the southern part of the Bridge. If the Bridge does contain two arms of gas from the SMC as suggested by previous authors, it is possible that an older population exists in the northern part of the Bridge that was tidally stripped as stars and not as gas or that formed stars in-situ. The results of the CMD analysis suggest that the Bridge may contain:

1. upper RGB stars to the tip of the RGB,
2. RGB and E-AGB stars,
3. Carbon stars, O-rich and dusty AGB stars,
4. reddened M-type giants.

These stars indicate ages older than previously suggested for the Bridge. The ages of the RGB and AGB stars in the central Bridge region is likely to range from $\sim 1-5$



Fig. 15. CMD, top is $(B - I)$ vs I of the blue objects of the Magellanic Bridge from SuperCOSMOS from one of the Bridge fields (α = 2:00:00, δ = −75:30:00) chosen in Sect. 4.2 prior to the removal of the galactic foreground population. Also shown are known (squares), and unknown (diamonds) blue objects in the northern part of the Bridge which remained after the removal of galactic foreground objects. Overlaid onto these populations are Padova isochrones (red lines) for stars at the estimated distance of the Magellanic Bridge (55 kpc with SMC metallicity (Z = 0.004) at a range of ages from log(age/yr) of 7.2 − 8.1 (top to bottom) with age steps of 0.1. Bottom is $(B - I)$ vs I of an offset field (α = 3:26:00, δ = −59:53:0) chosen in Sect. 4.2 for comparison to the Bridge population .

Gyr meaning that these stars were drawn into the Bridge at the formation event and did not form in-situ. It is possible from the observation of an older population that the northern part of the Bridge represents a different population from the southern arm not connected to the LMC and was formed from a different region of the SMC that contained stars at the time of the stripping event. Alternatively the Bridge may not be as young as previously thought. In order to better constrain the above, follow up observations of the Bridge populations are required as part of the VMC survey and also of metallicities and radial velocities from spectra that may constrain the membership of Bridge stars to the LMC or SMC.

Acknowledgements. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has made use of data from the DENIS public science archive as well as data obtained from the SuperCOSMOS Science Archive, prepared and hosted by the Wide Field Astronomy Unit, Institute for Astronomy, University of Edinburgh, which is funded by the UK Science and Technology Facilities Council and also the SIMBAD database, operated at CDS, Strasbourg, France.

References

Abraham, P., Dobashi, K., Mizuno, A., & Fukui, Y. 1995, A&A, 300, 525
Bekki, K. 2008, MNRAS, 390, L24
Bessell, M. S. & Brett, J. M. 1988, PASP, 100, 1134
Cannon, A. J. & Pickering, E. C. 1918, Annals of Harvard College Observatory, 91, 1
Chen, C.-H. R., Indebetouw, R., Chu, Y.-H., et al. 2010, ApJ, 721, 1206
Cioni, M.-R. L., Clementini, G., Girardi, L., et al. 2011, A&A, 527, A116+
Cioni, M.-R. L., Girardi, L., Marigo, P., & Habing, H. J. 2006, A&A, 452, 195
Cioni, M.-R. L., van der Marel, R. P., Loup, C., & Habing, H. J. 2000, A&A, 359, 601
Deinzer, W. & Schmidt, T. 1973, A&A, 27, 85
Demers, S. & Irwin, M. J. 1991, A&AS, 91, 171
Emerson, J. P., Sutherland, W. J., McPherson, A. M., et al. 2004, The Messenger, 117, 27
Epchtein, N. 1994, Experimental Astronomy, 3, 73
Frogel, J. A. & Blanco, V. M. 1990, ApJ, 365, 168
Girardi, L., Bressan, A., Bertelli, G., & Chiosi, C. 2000, A&AS, 141, 371
Grondin, L., Demers, S., & Kunkel, W. E. 1992, AJ, 103, 1234
Grondin, L., Demers, S., Kunkel, W. E., & Irwin, M. J. 1991, in IAU Symposium, Vol. 148, The Magellanic Clouds, ed. R. Haynes & D. Milne, 478
Hambly, N. C., Dufton, P. L., Keenan, F. P., et al. 1994, A&A, 285, 716
Hambly, N. C., MacGillivray, H. T., Read, M. A., et al. 2001, MNRAS, 326, 1279
Harris, J. 2007, ApJ, 658, 345
Harris, J. & Zaritsky, D. 2004, AJ, 127, 1531
Hindman, J. V., Kerr, F. J., & McGee, R. X. 1963a, Australian Journal of Physics, 16, 570
Hindman, J. V., McGee, R. X., Carter, A. W. L., Holmes, E. C. J., & Beard, M. 1963b, Australian Journal of Physics, 16, 552
Hog, E., Kuzmin, A., Bastian, U., et al. 1998, A&A, 335, L65
Holmberg, E. B., Lauberts, A., Schuster, H.-E., & West, R. M. 1974, A&AS, 18, 463
Iben, Jr., I. & Renzini, A. 1982, JRASC, 76, 398
Irwin, M. J., Demers, S., & Kunkel, W. E. 1990, AJ, 99, 191
Kennicutt, Jr., R. C. 1989, ApJ, 344, 685
Kerr, F. J. & de Vaucouleurs, G. 1955, Australian Journal of Physics, 8, 508
Mastropietro, C., Moore, B., Mayer, L., Stadel, J., & Wadsley, J. 2004, in Astronomical Society of the Pacific Conference Series, Vol. 327, Satellites and Tidal Streams, ed. F. Prada, D. Martinez Delgado, & T. J. Mahoney, 234
Mathewson, D. S. 1985, Proceedings of the Astronomical Society of Australia, 6, 104
McGee, R. X. & Newton, L. M. 1981, Proceedings of the Astronomical Society of Australia, 4, 189
Muller, E. & Bekki, K. 2007, MNRAS, 381, L11
Muller, E., Stanimirović, S., Rosolowsky, E., & Staveley-Smith, L. 2004, ApJ, 616, 845
Muller, E., Staveley-Smith, L., Zealey, W., & Stanimirović, S. 2003, MNRAS, 339, 105
Nikolaev, S. & Weinberg, M. D. 2000, ApJ, 542, 804
Olsen, K. A. G., Zaritsky, D., Blum, R. D., Boyer, M. L., & Gordon, K. D. 2011, ArXiv e-prints
Pakzad, S. L., Majewski, S. R., Frinchaboy, P. M., et al. 2004, in Bulletin of the American Astronomical Society, Vol. 36, American Astronomical Society Meeting Abstracts, 142.05
Putman, M. E. & Gibson, B. K. 1999, in Astronomical Society of the Pacific Conference Series, Vol. 166, Stromlo Workshop on High-Velocity Clouds, ed. B. K. Gibson & M. E. Putman, 276
Renzini, A. & Fusi Pecci, F. 1988, ARA&A, 26, 199
Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, AJ, 131, 1163
Smith, G. H. & Redenbaugh, A. K. 2010, PASP, 122, 1303
Westerlund, B. E. & Glaspey, J. 1971, A&A, 10, 1

Appendix A: Figures



Fig. A.1. A comparison of magnitude versus error for the (top to bottom) 2MASS J, H, K_s bands.



Fig. A.2. A comparison of magnitude versus error for the (top to bottom) DENIS I, J, K bands.